



Unrestricted					
Data and Business Rules – Atrial fibrillation Indicator Set					
Author	Paul Amos	Version No	14.0	Version Date	01-May-2009

New GMS Contract QOF Implementation
Dataset And Business Rules
-
Atrial Fibrillation

Amendment History:

Version	Date	Amendment History
		The version number starts at 3.1 in order to coincide with existing datasets and business rules.
Draft 3.1	21-Nov-2005	From Phil Brown
Draft 3.2	22-Nov-2005	Amended following review by Peter Horsfield
Draft 3.3	3-Dec-2005	Draft revised for internal review
7.4	23-Feb-2006	Amended following internal & 4 Countries review. Also includes uplifting to 7.4 to bring the 4-Byte documents to the same version number as the 5-Byte/CTV3/SNOMED-CT documents
8.0	15-Mar-2006	Signed off following 4 Country review
8.1	18-May-2006	Responding to queries raised Amend wording for Note 3
8.5	18-May-2006	Approved by NHSE
8.6	20-Oct-2006	April Read Code Release October Read Code Release Corrections and amendments following feedback
9.0	30-Nov-2006	Approved by NHSE
9.1	11-Apr-2007	April 2007 Read Code Release
9.2	11-June-2007	Following 4-Country Review: Changes to the qualifying criteria for AFIB_COD and AFIBDI_COD. Change date check for "anti-coagulant drug therapy or an anti-platelet drug therapy" (AF03) to 6 months
10.0	18-Jun-2007	Signed off following 4 Country review
10.1	23-Sep-2007	October 2007 Read Code Release
11.0	28-Nov-2007	Signed off following 4 Country review
11.1	30-Jun-2008	April 2008 Read Code Release QOF Review 2007 (Replacing AF2 with AF4)
12.0	24-Jul-2008	Signed off following 4 Country review
12.1	06-Oct-2008	October 2008 Read Code Release
13.0	05-Dec-2008	Signed off following 4 Country review
13.2	09-Mar-2009	QOF Review 2008
14.0	01-May-2009	Signed off following 4 Country review

New GMS contract Q&O framework implementation

Dataset and business rules – Atrial fibrillation indicator set

Notes

- 1) The specified dataset and rulesets are to support analysis of extracted data to reflect the status at a specified point in time of patient records held by the practice. In the context of this document that specified time point is designated the 'Reference date' and identified by the abbreviation 'REF_DAT'. In interpreting the specification REF_DAT should be taken to mean midnight of the preceding day (i.e. a REF_DAT of 01.04.2003 equates to midnight on 31.03.2003).
- 2) To support accurate determination of the population of patients to which the indicators should relate (the denominator population) these rulesets have been compiled with a prior assumption that the reference date is specified prior to extraction of data and is available for computation in the data extraction routine. The reference date will also be required to be included in the data extraction to support processing of rules that are dependent upon it. It is possible that an alternative approach could be adopted in which rules to determine the denominator population by registration status would be applied as a component of rule processing. If this second approach were to be adopted it would be essential to specify default time criteria for determining the registration characteristics of the denominator population during the data extraction process. Additionally there would be a requirement to supplement the dataset and rulesets to support identification of the appropriate denominator population.
- 3) Clinical codes quoted are (where known) from the October 2008 release of Read codes (version 0). The codes are shown within the document as a 4 character value to show that the Read Code is for a 4-Byte system.
 - i) Where a '%' wildcard is displayed, the Read Code is filled to 4 characters with full-stops. When implementing a search for the Read Code, only the non full-stop values should be used in the search, For example, a displayed Read Code of c1.% should be implemented as a search for c1%, i.e. should find c1 and any of it's children.
 - ii) Where a range of read codes are displayed, the Read Code is filled to 4 characters with full-stops. When implementing the search, only the non full-stop values should be used in the search, For example, a displayed Read Code range of G342 – G3z. should find all codes between G342 and G3z (including any children where applicable).

The version number starts at 3.1 in order to coincide with existing datasets and business rules

- 4) Datasets comprise a specification of two elements:
 - a) Patient selection criteria. These are the criteria used to determine the patient population against whom the indicators are to be applied.
 - i) Registration status. This determines the current patient population at the practice
 - ii) Diagnostic code status. This determines the current patient population (register size) for a given clinical condition

There are three scenarios within the diagnostic code status, these are where

- There is a single morbidity patient population (disease register) required (e.g. within CHD). Where this occurs, a single set of rules for identifying the patient population is provided.
- There is a single co-morbidity patient population (disease register) required (e.g. within Smoking). Where this occurs, a set of rules for **each** morbidity is provided. A patient **must** only be included in the patient population (register size) **once**.
- There are multiple patient populations (disease registers) required (e.g. within Heart Failure). Where this occurs, a single set of rules for **each** patient population is provided.

N.B. where there are multiple patient populations (disease registers), it is possible that one or more will also be a co-morbidity patient population (e.g. within Depression)

Where this occurs, details of which register population applies to which indicator(s) are provided. Where the register size applies to an indicator, this is the base denominator population for that indicator.

- b) Clinical data extraction criteria. These are the data items to be exported from the clinical system for subsequent processing to calculate points allocations. They are expressed in the form of a MIQUEST 'Report-style' extract of data.

The record of each patient that satisfies the appropriate selection criteria for a given indicator will be interrogated against the clinical data criteria (also appropriate to that indicator). A report of the data contained in the selected records will be exported in the form of a fixed-format tabular report. Each selected patient will be represented by a single row in the report. Rows will contain a fixed number of fields each containing a single data item. The number of fields in each row and their data content will be determined by the clinical data criteria. Data items that match the clinical data criteria will be exported in the relevant field of the report. Where there is no data to match a specific clinical criterion a null field will be exported.

- 5) Rulesets are specified as multiple rules to be processed sequentially. Processing of rules should terminate as soon as a 'Reject' or 'Select' condition is encountered
- 6) Rules are expressed as logical statements that evaluate as either 'true' or 'false'. The following operators are required to be supported:
- | | |
|---------------------|--------|
| a) > (greater than) | e) AND |
| b) < (less than) | f) OR |
| c) = (equal to) | g) NOT |
| d) ≠ (not equal to) | |
- 7) Where date criteria are specified with intervals of multiples of months or years these should be interpreted as calendar months or calendar years.
- 8) The new GMS contract requires that influenza vaccinations should be given between 1st September and 31st March of any given contract year in order to qualify for the relevant indicators. Hence in the contract year 2004 – 2005 the relevant dates will be 1st September 2004 and 31st March 2005 inclusive. In this document these dates are expressed as variable parameters FLU_COM and FLU_END respectively. For the purposes of data extraction these variables will be required to be specified prior to processing the relevant rules.

Dataset Specification**1) Patient selection criteria:**

a) Registration status

<i>Current registration status</i>	<i>Qualifying criteria</i>
Currently registered for GMS	Most recent registration date < (REF_DAT)
Previously registered for GMS	Any sequential pairing of registration date and deregistration date where both of the following conditions are met: registration date < (REF_DAT); and deregistration date >= (REF_DAT)

b) Diagnostic code status

<i>Code criteria</i>	<i>Qualifying diagnostic codes</i>	<i>Time criteria</i>
<i>Included</i>	<i>Read codes v0</i>	<i>Latest < (REF_DAT)</i>
	<i>G67.%</i>	
	<i>(Atrial fibrillation codes)</i>	
<i>Excluded</i>	<i>Read codes v0</i>	<i>Latest < (REF_DAT)</i> <i>AND > Date of</i> <i>diagnostic code above</i>
	<i>212R</i>	
	<i>(Atrial fibrillation resolved codes)</i>	

<u>Field Number</u>	<u>Field name</u>	<u>Data item</u>	<u>Qualifying criteria</u>
1	PAT_ID	Patient ID number	Unconditional
2	REG_DAT	Date of patient registration	Latest < (REF_DAT)
3	AFIBEXC_COD	<i>Read codes v0</i>	Latest < (REF_DAT)
		9hF1 9hF0	
		<i>(Atrial fibrillation exception reporting codes)</i>	
4	AFIBEXC_DAT	Date of AFIBEXC_COD	Chosen record
5	AFIB_COD	<i>Read codes v0</i>	Earliest < (REF_DAT)
		G67.%	
		<i>(Atrial fibrillation codes)</i>	
6	AFIB_DAT	Date of AFIB_COD	Chosen record
7	AFIBDI_COD	<i>Read codes v0</i>	Earliest < (REF_DAT) AND >= (AFIB_COD - 3 months)
		3272 8H41 8H44 8H4R 8HR1 8HVJ	
		<i>(Atrial fibrillation diagnosis codes)</i>	
8	AFIBDI_DAT	Date of AFIBDI_COD	Chosen record
9	XSAL_COD	<i>Read codes v0</i>	Latest < REF_DAT
		14LK Q545	
		<i>(Salicylate contra-indications: persistent)</i>	
10	XSAL_DAT	Date of XSAL_COD	Chosen record
11	TXSAL_COD	<i>Read codes v0</i>	Latest < REF_DAT
		8I24 8I38 8I66 8I70	
		<i>(Salicylate contra-indications: expiring)</i>	
12	TXSAL_DAT	Date of TXSAL_COD	Chosen record
13	XWAR_COD	<i>Read codes v0</i>	Latest < REF_DAT
		14LP Q565	

		<i>(Warfarin contraindications: persistent)</i>	
14	XWAR_DAT	Date of XWAR_COD	Chosen record
15	TXWAR_COD	<i>Read codes v0</i>	Latest < REF_DAT
		8I25 8I3E 8I65 8I71 8I2R 8I3d 8I6N 8I7A	
		<i>(Warfarin contraindications: expiring)</i>	
16	TXWAR_DAT	Date of TXWAR_COD	Chosen record
17	XCLO_COD	<i>Read codes v0</i>	Latest < REF_DAT
		14LQ Q566	
		<i>(Clopidogrel contraindications: persistent)</i>	
18	XCLO_DAT	Date of XCLO_COD	Chosen record
19	TXCLO_COD	<i>Read codes v0</i>	Latest < REF_DAT
		8I2K 8I3R 8I6B 8I72	
		<i>(Clopidogrel contraindications: expiring)</i>	
20	TXCLO_DAT	Date of TXCLO_COD	Chosen record
21	OSAL_COD	<i>Read codes v0</i>	Latest < REF_DAT
		67I8 8B63 8B3T 8B6P	
		<i>(OTC salicylate codes)</i>	
22	OSAL_DAT	Date of OSAL_COD	Chosen record
23	SAL_COD	<i>Read codes v0</i>	Latest < REF_DAT
		bu2.% di1.% j11.% blm.% bu4.%	
		<i>(Salicylate prescription codes)</i>	
24	SAL_DAT	Date of SAL_COD	Chosen record

25	CLO_COD	<i>Read codes v0</i>	Latest < REF_DAT
		bu5.%	
		<i>(Clopidogrel prescription codes)</i>	
26	CLO_DAT	Date of CLO_COD	Chosen record
27	WAR_COD	<i>Read codes v0</i>	Latest < REF_DAT
		bs..% 8B2K	
		<i>(Warfarin prescription codes)</i>	
28	WAR_DAT	Date of WAR_COD	Chosen record
29	DIPY_COD	<i>Read codes v0</i>	Latest < REF_DAT
		bu1.- bu12 bu14- bu1E	
		<i>(Dipyridamole prescription codes)</i>	
30	DIPY_DAT	Date of DIPY_COD	Chosen record
31	XDIPY_COD	<i>Read codes v0</i>	Latest < REF_DAT
		Q568 14LX	
		<i>(Dipyridamole contraindications: persistent)</i>	
32	XDIPY_DAT	Date of XDIPY_COD	Chosen record
33	TXDIPY_COD	<i>Read codes v0</i>	Latest < REF_DAT
		8I6a 8I7J 8I2b 8I3n	
		<i>(Dipyridamole contraindications: expiring)</i>	
34	TXDIPY_DAT	Date of TXDIPY_COD	Chosen record

Indicator rulesets

- 1 Indicator AF 1: The practice can produce a register of patients with Atrial Fibrillation

The terms of this indicator will be satisfied if the practice is able to produce a data extraction according to the above criteria.

No numerator or denominator determination is required.

- 2 Indicator AF 4: The percentage of patients with atrial fibrillation diagnosed from 1st April 2008 with ECG or specialist confirmed diagnosis

a) Denominator ruleset

<i>Rule number</i>	<i>Rule</i>	<i>Action if true</i>	<i>Action if false</i>
1	If AFIB_DAT >= 01.04.2008	Next rule	Reject
2	If AFIBDI_DAT <= (AFIB_DAT + 3 months) AND If AFIBDI_DAT >= (AFIB_DAT - 3 months)	Select	Next rule
3	If REG_DAT >= (REF_DAT - 3 months)	Reject	Next rule
4	If AFIBEXC_DAT >= (REF_DAT - 15 months)	Reject	Next rule
5	If AFIB_DAT >= (REF_DAT - 3 months)	Reject	Select

b) Numerator ruleset: To be applied to the above denominator population

<i>Rule number</i>	<i>Rule</i>	<i>Action if true</i>	<i>Action if false</i>
1	If AFIBDI_DAT <= (AFIB_DAT + 3 months) AND If AFIBDI_DAT >= (AFIB_DAT - 3 months)	Select	Reject

- 3 Indicator AF 3: The percentage of patients with atrial fibrillation who are currently treated with anti-coagulant drug therapy or an anti-platelet drug therapy

c) Denominator ruleset

<i>Rule number</i>	<i>Rule</i>	<i>Action if true</i>	<i>Action if false</i>
1	If SAL_DAT >= (REF_DAT - 6 months) OR If WAR_DAT >= (REF_DAT - 6 months) OR If CLO_DAT >= (REF_DAT - 6 months) OR If OSAL_DAT >= (REF_DAT - 6 months) OR If DIPY_DAT >= (REF_DAT - 6 months)	Select	Next rule
2	If REG_DAT >= (REF_DAT - 3 months)	Reject	Next rule
3	If AFIBEXC_DAT >= (REF_DAT - 15 months)	Reject	Next rule
4	If AFIB_DAT < (REF_DAT - 3 months)	Next rule	Reject
5	If XSAL_COD = Null AND If TXSAL_DAT = Null	Select	Next rule
6	If XSAL_COD = Null AND If TXSAL_DAT < (REF_DAT - 15 months)	Select	Next rule
7	If XWAR_COD = Null AND If TXWAR_DAT = Null	Select	Next rule
8	If XWAR_COD = Null AND If TXWAR_DAT < (REF_DAT - 15 months)	Select	Next rule
9	If XCLO_COD = Null AND If TXCLO_DAT = Null	Select	Next rule
10	If XCLO_COD = Null AND If TXCLO_DAT < (REF_DAT - 15 months)	Select	Next rule
11	If XDIPY_COD = Null AND If TXDIPY_DAT = Null	Select	Next rule
12	If XDIPY_COD = Null AND If TXDIPY_DAT < (REF_DAT - 15 months)	Select	Reject

d) Numerator ruleset: To be applied to the above denominator population

<i>Rule number</i>	<i>Rule</i>	<i>Action if true</i>	<i>Action if false</i>
1	If SAL_DAT >= (REF_DAT - 6 months) OR If WAR_DAT >= (REF_DAT - 6 months) OR If CLO_DAT >= (REF_DAT - 6 months) OR If OSAL_DAT >= (REF_DAT - 6 months) OR If DIPY_DAT >= (REF_DAT - 6 months)	Select	Reject